

CAUSAL PERTURBATION THEORY AND DIFFERENTIAL RENORMALIZATION

DIRK PRANGE

II. Institut für Theoretische Physik
Universität Hamburg
Luruper Chaussee 149
D-22761 Hamburg, Germany
`dirk.prange@desy.de`

ABSTRACT. In Causal Perturbation Theory the process of renormalization is precisely equivalent to the extension of time ordered distributions to coincident points. This is achieved by a modified Taylor subtraction on the corresponding test functions. I show that the pullback of this operation to the distributions yields expressions known from Differential Renormalization. The subtraction is equivalent to BPHZ subtraction in momentum space. Some examples from Euclidean scalar field theory in flat and curved spacetime will be presented.

INTRODUCTION

Calculations in perturbative QFT are performed primarily in momentum space. The computation of a given contribution to the S -Matrix is done by writing down Feynman rules and applying a certain choice of renormalization scheme to the resulting expression. For a reasonable renormalization scheme it should be proved to work to all orders and so produce a finite S -Matrix, as for example in the case of BPHZ renormalization.

But today we consider the principle of locality to be of special importance, and hence a local formulation of perturbation theory should exist. Indeed this was elaborated upon by Epstein and Glaser [EG73] following earlier ideas of Bogoliubov [BS76]. Their approach is called Causal Perturbation Theory (CPT). Based on a set of axioms they constructed the S -Matrix as a formal power series inductively. The process of renormalization occurs only once in every step. All lower order contributions are already renormalized. This corresponds to the determination of all divergent subgraphs in the traditional approach and simplifies the proof of the construction to all orders. The main concept on which CPT is based is its formulation completely in configuration space. During the seventies there have not been many applications of it except in the works [BS75, DM75]. This may be due to the fact that Epstein and Glaser used rigorous functional analysis, so renormalization is defined by an appropriate subtraction on test functions, whereas physicists are used to working with distributions in an integral kernel representation.

Later, Scharf et al. applied CPT to QED (see [Sch95] and references therein) and to non Abelian gauge theories [DHKS94a, DHKS94b, DHKS95, DHS95]. But their perturbative calculations are still performed in momentum space. In my opinion CPT has much potential for further applications to computation of diagrams in configuration space.

On the other hand, a renormalization scheme called Differential Renormalization [FJL92, SZ93] has gained attention. It is appreciated for its simplicity especially since no regularization procedure is needed. Differential Renormalization works in

configuration space on a large number of examples, but a proof for all orders is missing.

In the present paper I will show that the integral kernel representation of the Epstein-Glaser renormalization procedure exactly yields Differential Renormalization. It will lead to a simple formula for the computation of diagrams. I will apply it to some examples from Euclidean scalar field theory and use the results for renormalization group computations.

CPT mainly relies on the principles of causality, translation invariance and the singularity structure of the Feynman propagator. Brunetti and Fredenhagen [BF97] implemented CPT on a globally hyperbolic spacetime by giving a local generalization of translation invariance. Here the local causality structure is preserved and the Feynman propagator is known to have Hadamard form. I will show how the corresponding distributions can be renormalized in the Euclidean case. This is achieved by an appropriate translation of their representations from flat spacetime to curved spacetime.

1. THE EXTENSION OF DISTRIBUTIONS

Following [Sto93, Fre96] it turns out that renormalization in CPT actually is an extension of distributions from the subspace of test functions whose support does not contain the origin to the space of all test functions. To treat the most general solution of that problem we will be concerned with the space of distributions $\mathcal{D}'(\mathbb{R}^n)$, the dual of $\mathcal{D}(\mathbb{R}^n)$, the space of test functions with compact support. Let $\alpha = \{\alpha_1, \dots, \alpha_n\} \in \mathbb{N}^n$ be a multi-index, we set $|\alpha| = \sum_{i=1}^n \alpha_i$ and $\alpha! = \prod_{i=1}^n \alpha_i!$.

$$\partial^\alpha = \frac{\partial^{|\alpha|}}{\partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}} \quad (1)$$

is a partial differential operator of order $|\alpha|$.

Remark. Note that all operations on distributions like differentiation and transformations of their arguments are defined by the corresponding operations on test functions.

This fact is referred to as “*in the sense of distributions*”. Writing

$$T(\varphi) = \int d^n x T(x) \varphi(x), \quad T \in \mathcal{D}'(\mathbb{R}^n), \quad \varphi \in \mathcal{D}(\mathbb{R}^n), \quad (2)$$

we call $T(x)$ integral kernel of T . Let $\mathcal{D}(\mathbb{R}^n \setminus \{0\}) = \{\varphi \in \mathcal{D}(\mathbb{R}^n) | 0 \notin \text{supp}(\varphi)\}$ denote the subspace of test functions whose support does not contain the origin and $\mathcal{D}'(\mathbb{R}^n \setminus \{0\})$ its dual.¹ Now we state the

Problem. *Given a distribution ${}^0T \in \mathcal{D}'(\mathbb{R}^n \setminus \{0\})$, how can we construct an extension $T \in \mathcal{D}'(\mathbb{R}^n)$, such that ${}^0T(\varphi) = T(\varphi)$ for $\varphi \in \mathcal{D}(\mathbb{R}^n \setminus \{0\})$?*

The solution of this problem requires the introduction of a quantity that measures the singularity of the distribution at the origin [Ste71].

Definition 1. A distribution $T \in \mathcal{D}'(\mathbb{R}^n)$ has scaling degree s at $x = 0$, if

$$s = \inf \{s' \in \mathbb{R} | \lambda^{s'} T(\lambda x) \xrightarrow{\lambda \rightarrow 0} 0 \text{ in the sense of distributions} \}. \quad (3)$$

¹The existence of the extension is guaranteed by the Hahn-Banach theorem. A solution for homogenous distributions can be found in [Hör90][Chap. III.2]

Let s be denoted by $\text{scal deg}(T)$, and define $\text{sing ord}(T) := [s] - n$, the singular order.²

The definition also holds if $T \in \mathcal{D}'(\mathbb{R}^n \setminus \{0\})$. Take the δ -distribution as an

Example 1. For $\delta \in \mathcal{D}'(\mathbb{R}^n)$ one has: $\delta(\lambda x) = |\lambda|^{-n} \delta(x)$. The scaling degree of δ is n , the singular order is zero.

The scaling degree of some special compositions of distributions can be computed quite easily. We state

Proposition 1. Let $T \in \mathcal{D}'(\mathbb{R}^n)$ or $\mathcal{D}'(\mathbb{R}^n \setminus \{0\})$, $\text{scal deg}(T) = s$ and β be a multi-index.

- I. $\text{scal deg}(x^\beta T) = s - |\beta|$.
- II. $\text{scal deg}(\partial^\beta T) = s + |\beta|$.
- III. $\text{scal deg}(w) \leq 0, \text{scal deg}(wT) \leq s, w \in \mathcal{D}(\mathbb{R}^n)$.
- IV. $\text{scal deg}(T_1 \otimes T_2) = s_1 + s_2$, if $\text{scal deg}(T_i) = s_i$, $i = 1, 2$.

The proof is skipped, we only note that all statements follow directly from the translation of the words “in the sense of distributions” and the use of the Banach-Steinhaus theorem (principle of uniform boundedness, applied to distributions) on point III.

Example 2. The scaling degree of $\delta^{(\alpha)} \in \mathcal{D}'(\mathbb{R}^n)$ is $|\alpha| + n$. The singular order is $|\alpha|$.

The solution of the problem depends on the sign of the singular order. Let us consider the simple case first.

Theorem 2. Let ${}^0T \in \mathcal{D}'(\mathbb{R}^n \setminus \{0\})$ with scaling degree $s < n$. Then there exists a unique $T \in \mathcal{D}'(\mathbb{R}^n)$ with scaling degree s and $T(\varphi) = {}^0T(\varphi)$ for all $\varphi \in \mathcal{D}(\mathbb{R}^n \setminus \{0\})$.

The proof can be found in [BF97]. If the scaling degree is not smaller than the space dimension, the singular order ω is zero or positive. In that case theorem 2 guarantees a unique extension on test functions that vanish at the origin up to order ω . Thus a general extension can be defined after performing a projection into that subspace. This is achieved by a kind of modified Taylor subtraction, called the

W-operation. Let $\mathcal{D}^\omega(\mathbb{R}^n)$ be the subspace of test functions vanishing up to order ω at 0. Define

$$\begin{aligned} W_{(\omega;w)} : \mathcal{D}(\mathbb{R}^n) &\mapsto \mathcal{D}^\omega(\mathbb{R}^n), \quad \varphi \mapsto W_{(\omega;w)}\varphi, \\ (W_{(\omega;w)}\varphi)(x) &= \varphi(x) - w(x) \sum_{|\alpha| \leq \omega} \frac{x^\alpha}{\alpha!} \left(\partial^\alpha \frac{\varphi}{w} \right)(0), \end{aligned} \quad (4)$$

with $w \in \mathcal{D}(\mathbb{R}^n)$, $w(0) \neq 0$.

The action of $W_{(\omega;w)}$ on φ can be written as

$$(W_{(\omega;w)}\varphi)(x) = \sum_{|\beta| = \omega+1} x^\beta \varphi_\beta(x), \quad (5)$$

with $\varphi_\beta \in \mathcal{D}(\mathbb{R}^n)$. It has the nice property

$$W_{(\omega;w)}w\varphi = wW_{(\omega;1)}\varphi. \quad (6)$$

² $[s]$ is the largest integer that is smaller than or equal to s .

With $(\partial^\alpha x^\gamma)(0) = \gamma! \delta_\alpha^\gamma$ it follows for $|\gamma| \leq \omega$:

$$W_{(\omega;w)} w x^\gamma = w W_{(\omega;1)} x^\gamma \equiv 0. \quad (7)$$

Now we can discuss the general case.

Theorem 3. *Let ${}^0T \in \mathcal{D}'(\mathbb{R}^n \setminus \{0\})$ with scaling degree $s \geq n$. Given $w \in \mathcal{D}(\mathbb{R}^n)$ with $w(0) \neq 0$, a multi-index α , $|\alpha| \leq \omega$ and constants $C^\alpha \in \mathbb{C}$, then there is exactly one distribution $T' \in \mathcal{D}'(\mathbb{R}^n)$ with scaling degree s and following properties:*

- I. $\langle T', \varphi \rangle = \langle {}^0T, \varphi \rangle \quad \forall \varphi \in \mathcal{D}(\mathbb{R}^n \setminus \{0\}),$
- II. $\langle T', w x^\alpha \rangle = C^\alpha.$

T' is given by:

$$\langle T', \varphi \rangle = \langle T, W_{(\omega;w)} \varphi \rangle + \sum_{|\alpha| \leq \omega} \frac{C^\alpha}{\alpha!} \left(\partial^\alpha \frac{\varphi}{w} \right) (0). \quad (8)$$

Here T is the unique extension by theorem 2, $W_{(\omega;w)}$ is given by (4) and ω is the singular order of 0T .

The proof can be found in [BF97]. We see that in the case of non-negative singular order, the extension is not unique. It is fixed by a finite set of complex numbers C^α . Let us look at the next

Example 3. The n -th power of the scalar Feynman propagator $(i\Delta_F)^n(x) = \theta(x^0)\Delta_+^n(x) + \theta(-x^0)\Delta_+^n(-x)$ is a distribution on $\mathbb{R}^4 \setminus \{0\}$. We compute the scaling degree of Δ_+^n .

$$\begin{aligned} \Delta_+^n(\lambda x) &= (2\pi)^{-3n} \int \prod_{i=1}^n \frac{d^3 \mathbf{p}_i}{2\omega_{\mathbf{p}_i}} e^{\sum_{i=1}^n (-i\omega_{\mathbf{p}_i} \lambda x^0 + i\mathbf{p}_i \lambda \mathbf{x})} \\ &= \lambda^{-2n} (2\pi)^{-3n} \int \prod_{i=1}^n \frac{d^3 \mathbf{p}_i}{2\sqrt{(\lambda m)^2 + \mathbf{p}_i^2}} e^{\sum_{i=1}^n (-i\sqrt{(\lambda m)^2 + \mathbf{p}_i^2} x^0 + i\mathbf{p}_i \mathbf{x})} \\ &= \lambda^{-2n} \Delta_+^n(x, \lambda m) \\ &\xrightarrow{\lambda \rightarrow 0} \lambda^{-2n} D_+^n(x). \end{aligned}$$

Here D_+ denotes the massless scalar two point function. Hence the scaling degree is $2n$. The application of the W -operation with $\omega = 2n - 4$ yields the extension to all test functions. The computation can be done similarly for the Euclidean propagator.

We turn to another example that seems to have caused some confusion in classical physics (see e.g. [FLS64]).

Example 4 (The self energy of the electron). In electrostatics the electric potential of an electron at the origin is given by the Green's function of the Laplace equation in 3 dimensions.

$$\Delta \phi = -4\pi \rho = 4\pi e \delta \quad \Rightarrow \quad \phi = -\frac{e}{r} \in \mathcal{D}'(\mathbb{R}^3).$$

The electric field is

$$\mathbf{E} = -\nabla \phi = -\frac{e\mathbf{r}}{r^3} \in \mathcal{D}'(\mathbb{R}^3).$$

Since $\text{sing supp}(\mathbf{E}) = \{0\}$ it follows:

$$\mathbf{E}^2 = \frac{e^2}{r^4} \in \mathcal{D}'(\mathbb{R}^3 \setminus \{0\}).$$

The singular order is 1. Hence there is an extension to all test functions by the W -operation. We can define the energy density $U = \mathbf{E}^2$ as the following distribution:

$$\langle U, \varphi \rangle := \langle \mathbf{E}^2, W_{(1;w)} \varphi \rangle. \quad (9)$$

An electron at rest has self energy $E = 1/(4\pi) \langle U, 1 \rangle$. The choice of $\varphi \equiv 1$ is possible due to sufficient convergence at long range. The same holds for w in (9).

$$E = \frac{1}{4\pi} \langle U, W_{(1;1)} 1 \rangle + C^0 = C^0,$$

as all C^α , $|\alpha| = 1$ vanish. We can determine C^0 by the requirement that the mass of the electron is purely electromagnetic, i.e.

$$E = mc^2.$$

In the following I will suppress the distinction between 0T and T in the case that the scaling degree is smaller than the space dimension. This should lead to no confusion since the extension is unique in that case.

1.1. The integral kernel representation. To use standard vocabulary we will call the extended distribution in theorem 3 a *renormalization*. Next we will work out its integral kernel. If we set all C^α to zero we have

Definition 2. Let $T \in \mathcal{D}'(\mathbb{R}^n)$ with $\text{sing ord}(T) = \omega$. The integral kernel $T_{R(\omega;w)}$ of its extension is given by

$$\langle T_{R(\omega;w)}, \varphi \rangle := \langle T, W_{(\omega;w)} \varphi \rangle. \quad (10)$$

Furthermore, we consider a family of distributions T_t that depend continuously on a real parameter t . If K is a real compact interval then $\int_K dt \langle T_t, \varphi \rangle$ exists as a Riemannian integral. We define

$$\left\langle \int_K dt T_t, \varphi \right\rangle := \int_K dt \langle T_t, \varphi \rangle \quad (11)$$

in the sense of distributions. Now we have

Proposition 4. *The integral kernel $(T_{R(\omega;w)} w)$ is given by:*

$$(T_{R(\omega;w)} w)(x) = (-)^{\omega+1} (\omega+1) \sum_{|\beta|=\omega+1} \frac{\partial^\beta x^\beta}{\beta!} \int_0^1 dt \frac{(1-t)^\omega}{t^{\omega+1}} T\left(\frac{x}{t}\right) w\left(\frac{x}{t}\right). \quad (12)$$

Proof. The Taylor expansion of φ at the origin is:

$$\varphi(x) = \sum_{|\alpha|=0}^{\omega} \frac{x^\alpha}{\alpha!} (\partial^\alpha \varphi)(0) + (\omega+1) \sum_{|\beta|=\omega+1} \frac{x^\beta}{\beta!} \int_0^1 dt (1-t)^\omega (\partial^\beta \varphi)(tx). \quad (13)$$

Hence $(W_{(\omega;1)}\varphi)(x)$ is the Taylor rest term of order $\omega+1$. Writing (12) as $(T_{R(\omega;w)}w) = \int_0^1 dt (T_{R(\omega;w)}w)_t$ we find using (11):

$$\begin{aligned} \left\langle \int_0^1 dt (T_{R(\omega;w)}w)_t, \varphi \right\rangle &= (\omega+1) \int_0^1 dt (1-t)^\omega \times \\ &\quad \times \int d^n x \sum_{|\beta|=\omega+1} \frac{x^\beta}{\beta!} T(x)w(x)(\partial^\beta \varphi)(tx) \\ &= \langle T, wW_{(\omega;1)}\varphi \rangle \\ &= \langle T, W_{(\omega;w)}w\varphi \rangle \\ &= \langle T_{R(\omega;w)}, w\varphi \rangle \\ &= \langle T_{R(\omega;w)}w, \varphi \rangle, \end{aligned}$$

where we used equation (6). $\square$

Note that the differential operator in (12) is a *weak* derivative. If w had no zeros ($\Rightarrow w \notin \mathcal{D}(\mathbb{R}^n)$), the integral kernel would result from a simple division. Later we will encounter a well known example of this.

To achieve a similar representation for the whole distribution we put a restriction on the test function w in (4). Let $w(0) = 1$ and $(\partial^\alpha w)(0) = 0$, for $|\alpha| \leq \omega$.

Remark. This is no real loss of generality since for a given $v \in \mathcal{D}(\mathbb{R}^n)$, $v(0) \neq 0$, the projection

$$v \mapsto w = v \sum_{|\alpha| \leq \omega} \frac{x^\alpha}{\alpha!} \left(\partial^\alpha \frac{1}{v} \right) (0) \in \mathcal{D}(\mathbb{R}^n) \quad (14)$$

yields $(\partial^\gamma w)(0) = \delta_0^\gamma$ for $|\gamma| \leq \omega$.

Then $(1-w)$ vanishes up to order ω at 0:

$$W_{(\omega;w)}(1-w)\varphi = (1-w)\varphi, \quad (15)$$

$$\langle (1-w)T_{R(\omega;w)}, \varphi \rangle = \langle (1-w)T, \varphi \rangle. \quad (16)$$

Now the integral kernel is given by

Lemma 5. *With the above restrictions on $w \in \mathcal{D}(\mathbb{R}^n)$, the renormalized distribution $T_{R(\omega;w)}$ has the following integral kernel:*

$$\begin{aligned} T_{R(\omega;w)}(x) &= (-)^\omega (\omega+1) \sum_{|\beta|=\omega+1} \partial^\beta \frac{x^\beta}{\beta!} \left[- \int_0^1 dt \frac{(1-t)^\omega}{t^{n+\omega+1}} T\left(\frac{x}{t}\right) w\left(\frac{x}{t}\right) + \right. \\ &\quad \left. + \int_1^\infty dt \frac{(1-t)^\omega}{t^{n+\omega+1}} T\left(\frac{x}{t}\right) (1-w)\left(\frac{x}{t}\right) \right]. \quad (17) \end{aligned}$$

Proof. The first term of (17) is the integral kernel of $wT_{R(\omega;w)}$. A simple computation shows that the second term smeared with φ yields $\langle T, (1-w)\varphi \rangle$. Equation (16) completes the proof. $\square$

If $n = 4m$, we can write

$$k! \sum_{|\beta|=k} \partial^\beta \frac{x^\beta}{\beta!} = \mathfrak{d} \left(\sum_{|\beta|=1} \partial^\beta x^\beta \right)^k \mathfrak{d} = \mathfrak{d} (\partial_{\mu_1} x^{\mu_1} + \dots + \partial_{\mu_m} x^{\mu_m})^k \mathfrak{d}, \quad (18)$$

$$\text{with } x = \begin{pmatrix} x^{\mu_1} \\ \vdots \\ x^{\mu_m} \end{pmatrix} \in \mathbb{R}^{4m}.$$

Here $\mathfrak{d} \dots \mathfrak{d}$ denotes the ordering of differential operators to the left of the coordinates.

To perform some computations with formula (17) it would be desirable to abandon the requirement of w being a test function. Let $w_m \in \mathcal{D}(\mathbb{R}^n)$ be a sequence with $\lim_{m \rightarrow \infty} w_m =: w \in \mathcal{D}'(\mathbb{R}^n)$. If $\lim_{m \rightarrow \infty} \langle T_{R(\omega; w_m)}, \varphi \rangle \in \mathbb{C}$ exists $\forall \varphi \in \mathcal{D}(\mathbb{R}^n)$, we will allow w to be used in the renormalization procedure. Let us consider $T \in \mathcal{D}^{\omega'}(\mathbb{R}^n)$ with singular order ω and $\text{sing supp}(T) = \{0\}$. Choose $w(x) = \theta(1/M - |x|) =: \theta^<(x)$, $M \in \mathbb{R}$, $M > 0$, where $|\cdot|$ denotes the Euclidean norm. Since $\text{sing supp}(T) \cap \text{sing supp}(\theta^<) = \emptyset$ and $\theta^<$ has compact support, the pointwise product $\theta^< T \in \mathcal{E}^{\omega'}(\mathbb{R}^n) \subset \mathcal{D}^{\omega'}(\mathbb{R}^n)$ exists.³ Applying (17) yields

$$T_{R(\omega; \theta^<)}(x) = (-)^\omega (\omega + 1) \sum_{|\beta|=\omega+1} \partial^\beta \frac{x^\beta}{\beta!} \int_1^{M|x|} dt \frac{(1-t)^\omega}{t^{n+\omega+1}} T\left(\frac{x}{t}\right) \quad (19)$$

$$=: T_R^M(x),$$

For an arbitrary choice of w , a scale M has to be introduced for dimensional reasons. This allows writing $w(Mx)$ as a function of a dimensionless argument. The dependence of the counterterms on the scale can easily be computed:

$$M \frac{\partial}{\partial M} \langle T_{R(\omega; w(Mx))}, \varphi \rangle = \sum_{|\alpha| \leq \omega} B^\alpha \langle \delta^{(\alpha)}, \varphi \rangle, \quad (20)$$

$$\text{with } B^\alpha = \frac{(-)^{|\alpha|+1}}{\alpha!} \langle T, M x^\mu (\partial_\mu w)(Mx) x^\alpha \rangle. \quad (21)$$

For $w = \theta^<$ we get

$$B^\alpha = \frac{(-)^{|\alpha|}}{\alpha!} \left\langle T, \delta \left(\frac{1}{M} - |x| \right) x^\alpha \right\rangle. \quad (22)$$

1.2. Momentum space and BPHZ renormalization. Since the Fourier transformation is a map $\mathcal{D} \mapsto \mathcal{S}$ we have to restrict our distributional space to $\mathcal{S}' \subset \mathcal{D}'$.⁴ We remind the reader that the Fourier transformation is defined in the sense of distributions, i.e. $\widehat{T}(\varphi) := T(\widehat{\varphi})$.

First we start with the definition of the moments of a test function:

$$K^\alpha(\psi) := \int d^n x x^\alpha \psi(x), \quad \psi \in \mathcal{S}(\mathbb{R}^n). \quad (23)$$

³ $\mathcal{E} = \mathcal{C}^\infty$ and $\mathcal{E}'$ is the space of distributions with compact support.

⁴Let $\mathcal{S}$ be the space of $\mathcal{C}^\infty$ functions of rapid decrease and $\mathcal{S}'$ its dual. We use the convention

$$\widehat{\psi}(p) = \int d^n x \psi(x) e^{ipx}.$$

Let $\mathcal{S}_\omega(\mathbb{R}^n) := \{\psi \in \mathcal{S}(\mathbb{R}^n), K^\alpha(\psi) = 0, |\alpha| \leq \omega\}$ be the subspace of test function with vanishing moments up to order ω , then it follows: $\psi \in \mathcal{S}^\omega \Rightarrow \check{\psi} \in \mathcal{S}_\omega$. Choosing w as in Lemma 5 we have:

$$K^\alpha(\widehat{w}) = 0 \text{ for } 0 < |\alpha| \leq \omega, \quad K^0(\widehat{w}) = (2\pi)^n, \quad (24)$$

and by a simple computation:

$$K^\gamma(\partial^\alpha \widehat{w}) = (-)^{|\gamma|} \gamma! \delta_\alpha^\gamma (2\pi)^n, \quad \gamma \leq \alpha. \quad (25)$$

The Fourier transformation of $W\varphi$ is:

$$(W_{(\omega;w)}\varphi)^\vee(p) = \check{\varphi}(p) - \frac{1}{(2\pi)^n} \sum_{|\alpha| \leq \omega} \frac{(-)^{|\alpha|}}{\alpha!} (\widehat{wx^\alpha})(-p) \langle \delta^{(\alpha)}, \varphi \rangle \quad (26)$$

$$= \check{\varphi}(p) - \frac{1}{(2\pi)^n} \sum_{|\alpha| \leq \omega} \frac{K^\alpha(\check{\varphi})}{\alpha!} (\partial^\alpha \widehat{w})(-p). \quad (27)$$

Using (25) we get:

$$K^\gamma \left((W_{(\omega;w)}\varphi)^\vee \right) = 0, \quad |\gamma| \leq \omega, \quad (28)$$

so $W_{(\omega;w)}^\vee$ actually is a projector $\mathcal{S} \mapsto \mathcal{S}_\omega$. With

$$\langle T_{R(\omega;w)}, \varphi \rangle = \langle \widehat{T_{R(\omega;w)}}, \check{\varphi} \rangle = \langle \widehat{T}, (W_{(\omega;w)}\varphi)^\vee \rangle \quad (29)$$

the integral kernel is given by

$$\widehat{T_{R(\omega;w)}}(k) = \widehat{T}(k) - \sum_{|\alpha|=0}^{\omega} \frac{k^\alpha}{\alpha!} (\partial^\alpha \widehat{T}w)(0). \quad (30)$$

This subtraction should be understood in the sense of distributions, i.e. the subtraction on the test functions has to occur before smearing out. It looks similar to BPHZ renormalization which is a Taylor subtraction at arbitrary momentum q . We will compute the corresponding w . Let

$$\widehat{T_R^q}(k) := \widehat{T}(k) - \sum_{|\alpha|=0}^{\omega} \frac{(k-q)^\alpha}{\alpha!} (\partial^\alpha \widehat{T})(q) \quad (31)$$

denote the BPHZ renormalized distribution in momentum space. Using the Taylor rest expression similar to (13) and performing Fourier transformation we get:

$$T_R^q(x) = (-)^{\omega+1} (\omega+1) e^{-iqx} \sum_{|\beta|=\omega+1} \partial^\beta \frac{x^\beta}{\beta!} \int_0^1 dt \frac{(1-t)^\omega}{t^{n+\omega+1}} T\left(\frac{x}{t}\right) e^{i\frac{qx}{t}}, \quad (32)$$

and comparing to (12),

$$= T_{R(\omega;e^{iqx})}. \quad (33)$$

Here, the equivalence of the subtraction procedure in Epstein-Glaser and BPHZ renormalization can be seen explicitly. Moreover this cannot be achieved by the requirement $w \in \mathcal{S}(\mathbb{R}^n)$.

2. CAUSAL PERTURBATION THEORY

In the following I will give a very brief summary of CPT. A complete description can be found in [EG73, Sch95].

We start with the S -Matrix as a formal power series

$$S(g) = 1 + \sum_{n=1}^{\infty} \frac{(-i)^n}{n!} \int d^4x_1 \cdots d^4x_n T_n(x_1, \dots, x_n) g(x_1) \cdots g(x_n). \quad (34)$$

It is an operator valued functional. The function $g \in \mathcal{D}(\mathbb{R}^4)$ plays the role of a coupling “constant”. The T_n are operator valued distributions in Fock space. They are called time ordered functions and involve free fields only. Epstein and Glaser stated six axioms from which the time ordered functions can be computed recursively. Here we cite only the most important one called the causality axiom:

$$T_n(x_1, \dots, x_n) = T_k(x_1, \dots, x_k) T_{n-k}(x_{k+1}, \dots, x_n), \quad (35)$$

if all points $x_{k+1}, \dots, x_n$ are not in the causal past of $x_1, \dots, x_k$. The inductive construction starts with $T_1 = \mathcal{L}_{\text{int}}$. We assume that $T_{n'}(x_1, \dots, x_{n'})$ for all $n' < n$ exist as a sum of products of a symmetric translation invariant numerical distribution and a Wick polynomial of fields. The scaling degree of the numerical distribution is known at coincident points. Now T_n can be constructed up to the total diagonal $x_1 = \dots = x_n$ by the causality axiom. Wick’s theorem ensures the required form. The numerical distributions have an extension to the diagonal which is the origin in $\mathbb{R}^{4n-4}$ because of translation invariance. The Wick polynomials are already defined as operator valued distributions on the whole space. Let me emphasize that the translation invariance of the numerical distributions plays a crucial role in the whole construction.

3. APPLICATIONS

I will give some examples from ϕ^4 -theory in lowest order. The Lagrangian for the self interacting scalar field is

$$\mathcal{L}(x) = \frac{1}{2} : \partial_\mu \phi(x) \partial^\mu \phi(x) : - \frac{m^2}{2} : \phi^2(x) : - \frac{\lambda}{4!} : \phi^4(x) : . \quad (36)$$

Now T_1 is given by the interaction term.

$$T_1(x) = -\frac{\lambda}{4!} : \phi^4(x) : . \quad (37)$$

Causality (35) implies T_2 for non coincident points.

$$\begin{aligned} T_2(x_1, x_2) &= \begin{cases} T_1(x_1)T_1(x_2), & \text{if } x_1^0 > x_2^0, \\ T_1(x_2)T_1(x_1), & \text{if } x_2^0 > x_1^0, \end{cases} \\ &= \frac{\lambda^2}{(4!)^2} \left[: \phi^4(x_1) \phi^4(x_2) : + \right. \end{aligned} \quad (38)$$

$$+ 16 \Delta_F(x_1 - x_2) : \phi^3(x_1) \phi^3(x_2) : + \quad (39)$$

$$+ 72 (i \Delta_F)^2(x_1 - x_2) : \phi^2(x_1) \phi^2(x_2) : + \quad (40)$$

$$+ 96 (i \Delta_F)^3(x_1 - x_2) : \phi(x_1) \phi(x_2) : + \quad (41)$$

$$\left. + 24 (i \Delta_F)^4(x_1 - x_2) \right]. \quad (42)$$

To give some explicit results for the numerical distributions we will now turn to their corresponding Euclidean counterparts. The singular support of these distributions is the origin only. Hence we can use (19) for the renormalization.

3.1. The massless theory. The Green's function of the Laplace equation in four dimensions is

$$D_F(x) = \frac{1}{4\pi^2} \frac{1}{x^2}. \quad (43)$$

It has singular order -2 . Consider the contribution to the two point resp. four point vertex function.



3.1.1. *The one-loop graph.* The “fish” graph (40) has singular order zero, so we get

$$D_F^2|_R^M(x) = \frac{1}{(2\pi)^4} \frac{1}{2} \partial_\mu x^\mu \frac{\ln(M^2 x^2)}{(x^2)^2}. \quad (44)$$

Since all extensions only differ by a δ -term we have

$$M \frac{\partial}{\partial M} D_F^2|_R^M(x) = \frac{1}{8\pi^2} \delta(x) \quad \text{or} \quad \partial_\mu \frac{x^\mu}{(x^2)^2} = 2\pi^2 \delta(x), \quad (45)$$

by (22). This also could have been seen by expressing $\frac{x^\mu}{(x^2)^2} = -\frac{1}{2} \partial^\mu \frac{1}{x^2}$, which is unique by theorem 2.

3.1.2. *The two-loop graph.* The singular order of the “setting sun” (41) is 2. Hence we have



$$D_F^3|_R^M(x) = \frac{1}{(2\pi)^6} \frac{1}{4} \partial_\mu \partial_\nu \partial_\sigma x^\mu x^\nu x^\sigma \frac{1}{(x^2)^3} \left[\ln(M^2 x^2) + M^2 x^2 - 4M\sqrt{x^2} + 3 \right]. \quad (46)$$

Terms two, three and four have singular order 0, 1 and 2. As they are zero outside the origin, they must be proportional to δ and its first resp. second derivatives. As there is no Euclidean invariant combination of ∂ and δ , term two has to be zero. Using (22) we get by comparison with $M \frac{\partial}{\partial M}$ on (46):

$$\partial_\mu \partial_\nu \partial_\sigma x^\mu x^\nu x^\sigma \frac{1}{(x^2)^3} = \frac{\pi^2}{2} \Delta \delta(x), \quad (47)$$

$$\partial_\mu \partial_\nu \partial_\sigma x^\mu x^\nu x^\sigma \frac{1}{(x^2)^2} = 4\pi^2 \delta(x), \quad (48)$$

$$\partial_\mu \partial_\nu \partial_\sigma x^\mu x^\nu x^\sigma \frac{1}{\sqrt{x^2}^5} \equiv 0, \quad (49)$$

where Δ is the Laplacian. Now we will turn to


3.2. The massive theory. The Green's function of the Euclidean Klein-Gordon equation is

$$\Delta_F(x) = \frac{1}{(2\pi)^2} \frac{m K_1(m\sqrt{x^2})}{\sqrt{x^2}}. \quad (50)$$

As $K_1(x) \propto 1/x$ for $x \rightarrow 0$, the singular order of $\Delta_F = -2$. We will compute

3.2.1. *The one-loop graph.* With $\text{sing ord}(\Delta_F^2) = 0$ and (19) we get:

$$(\Delta_F^2)|_R^M(x) = \frac{1}{32\pi^4} \partial_\mu x^\mu \left\{ \frac{m^2}{x^2} \left[K_1^2(m\sqrt{x^2}) - K_0(m\sqrt{x^2}) K_2(m\sqrt{x^2}) \right] + \right. \\ \left. - \frac{m^2}{M^2} \frac{1}{(x^2)^2} \left[K_1^2\left(\frac{m}{M}\right) - K_0\left(\frac{m}{M}\right) K_2\left(\frac{m}{M}\right) \right] \right\}. \quad (51)$$



By writing $x^\mu \dots = \partial^\mu \dots$ which is unique we can express it as:

$$= \frac{m^2}{2(2\pi)^4} \Delta \left\{ K_0^2(m\sqrt{x^2}) - K_1^2(m\sqrt{x^2}) + \frac{K_0(m\sqrt{x^2}) K_1(m\sqrt{x^2})}{m\sqrt{x^2}} + \right. \\ \left. + \frac{1}{2M^2 x^2} \left[K_1^2\left(\frac{m}{M}\right) - K_0\left(\frac{m}{M}\right) K_2\left(\frac{m}{M}\right) \right] \right\}. \quad (52)$$

This can be compared to the corresponding expression in [HL92].

3.2.2. *The two-loop graph.* Applying formula (19) leads to an integral $\int ds (s - \text{const})^2 K_1^3(s)$ that is difficult to solve. But if we use the expansion

$$\frac{m^3 K_1^3(mr)}{r^3} = \frac{1}{r^6} + \frac{3m^2}{4r^4} \left(\ln\left(\frac{m^2 r^2}{4}\right) + \ln(\gamma^2) - 1 \right) + R(r^{-2}), \quad (53)$$

we can renormalize the first and second summand with singular order 2, 0 respectively. With

$$\left. \frac{\ln\left(\frac{m^2 x^2}{4}\right)}{(x^2)^2} \right|_R^M = \frac{1}{4} \partial_\mu x^\mu \frac{\ln\left(\frac{m^4 x^2}{16M^2}\right) \ln(M^2 x^2)}{(x^2)^2} \quad (54)$$

and the previous results from subsection 3.1 we get:

$$(\Delta_F^3)|_{MR}^M(x) = \frac{1}{(2\pi)^6} \left\{ \frac{1}{4} \partial_\mu \partial_\nu \partial_\sigma x^\mu x^\nu x^\sigma \frac{1}{(x^2)^3} [\ln(M^2 x^2) + M^2 x^2 + 3] + \right. \\ \left. + \frac{3m^2}{16} \partial_\mu x^\mu \frac{1}{(x^2)^2} \left[\ln\left(\frac{m^4 x^2}{16M^2}\right) + 2 \ln(\gamma^2) - 2 \right] \ln(M^2 x^2) + R(x^{-2}) \right\}, \quad (55)$$



where the subscript MR denotes the “minimal renormalization”, i.e. every summand is renormalized with its singular order.

3.3. The renormalization group. The parameter M plays the role of a renormalization scale. It enters the theory by purely dimensional reasons as an argument of the “function” w . We will give the lowest order contributions to the β , γ and γ_m functions in the renormalization group. They can be read off by solving the renormalization group equation

$$\left[M \frac{\partial}{\partial M} + \beta(g) \frac{\partial}{\partial g} + m \gamma_m(g) \frac{\partial}{\partial m} - n \gamma(g) \right] \Gamma_n(x_1, \dots, x_n) = 0 \quad (56)$$

to order g^2 , where $g = \lambda/(16\pi^2)$. Using the expansions

$$\beta(g, m, M) = M \frac{\partial g}{\partial M} = \sum_{n=2}^{\infty} \beta_n g^n, \quad (57)$$

$$\gamma_m(g, m, M) = M \frac{\partial \ln m}{\partial M} = \sum_{n=2}^{\infty} \gamma_{m,n} g^n, \quad (58)$$

$$\gamma(g, m, M) = \frac{M}{2} \frac{\partial \ln Z}{\partial M} = \sum_{n=2}^{\infty} \gamma_n g^n, \quad (59)$$

we find $\beta_2 = 3$ and $\gamma_2 = \frac{1}{12}$ for the massless theory. Here we had to add a term to the setting sun proportional to (48) to achieve $\gamma_m \equiv 0$.

In the massive theory we find

$$\beta_2 = 3 \frac{m^2}{M^2} K_1^2 \left(\frac{m}{M} \right), \quad (60)$$

hence $\beta_2 \rightarrow 3$ if $M \rightarrow \infty$. This result also holds for the corresponding minimal renormalization. In that scheme we get

$$\gamma_2 = \frac{1}{12}, \quad (61)$$

$$\gamma_{m,2} = \frac{2M^2}{3m^2} + \frac{1}{2} \ln \left(\frac{m^2}{4M^2} \right) + \ln(\gamma) - \frac{5}{12}, \quad (62)$$

by using (55).

3.4. Curved spacetime. Let $\mathcal{M}$ be a globally hyperbolic manifold with a metric g . Wick polynomials were defined in [BFK96] using techniques from microlocal analysis. Then CPT was implemented in [BF97] for scalar ϕ^4 -theory. The Feynman propagator is known to have Hadamard structure [KW91]:

$$\Delta_F \propto \frac{\Delta^{\frac{1}{2}}}{2\sigma} + v \ln(2\sigma) + w, \quad (63)$$

where σ, Δ, v , and w are smooth functions on the manifold and an appropriate $i\epsilon$ regularization has to be chosen. By using a chart it can be seen to have the same scaling degree as in flat spacetime. The function σ is half the square of the geodesic distance which is unique in every sufficiently small neighbourhood. Let $g = \det(g_{ab})$. The Van-Vleck-Morette determinant

$$\Delta(p, p') = -\frac{1}{\sqrt{g(p)g(p')}} \det(-\sigma_{ab'}(p, p')) \quad (64)$$

fulfills the following differential equation:

$$\nabla_a (\Delta \sigma^a) = 4\Delta. \quad (65)$$

The vector index on σ denotes the covariant derivative as usual. If we expand

$$v = \sum_{n=0}^{\infty} v_n \sigma^n, \quad w = \sum_{n=0}^{\infty} w_n \sigma^n, \quad (66)$$


in powers of σ , the coefficients (except w_0) are determined by the Hadamard recursion relations, see [DB60].

Following [BF97] the causal construction of the second order S -Matrix is the same as in flat spacetime. The only step left is to perform the extension to coincident points.

Therefore we turn to the corresponding Euclidean distributions, so we can transfer our results from the previous examples.⁵ They may be compared with [Lüs82, Bun81]. Again we only use the minimal renormalization scheme.

3.4.1. *The one-loop graph.* This is given by

$$\begin{aligned} \Delta_F^2|_{MR}^M &= \frac{1}{16\pi^4} \left(\frac{\Delta}{(2\sigma)^2} \Big|_R^M + 2 \frac{\Delta^{\frac{1}{2}} v \ln(2\sigma)}{2\sigma} + 2 \frac{\Delta^{\frac{1}{2}} w}{2\sigma} + 2vw \ln(2\sigma) + v^2 \ln^2(2\sigma) + w^2 \right). \end{aligned} \quad (67)$$



Only the first term needs renormalization and we get

$$\frac{\Delta}{(2\sigma)^2} \Big|_R^M = \frac{1}{2} \nabla_a \sigma^a \frac{\Delta \ln(2M^2 \sigma)}{(2\sigma)^2}. \quad (68)$$

3.4.2. *The two-loop graph.* Here we use the expansion (66) to determine the terms that have to be renormalized. Then we have

$$\begin{aligned} \Delta_F^3|_{MR}^M &= \frac{1}{(2\pi)^6} \left(\frac{\Delta^{\frac{3}{2}}}{(2\sigma)^3} \Big|_R^M + 3 \frac{\Delta v_0 \ln(2\sigma)}{(2\sigma)^2} \Big|_R^M + 3 \frac{\Delta w_0}{(2\sigma)^2} \Big|_R^M + 3 \frac{\Delta \bar{v} \ln(2\sigma)}{2\sigma} + \right. \\ &\quad + 3 \frac{\Delta \bar{w}}{2\sigma} + 3 \frac{\Delta^{\frac{1}{2}} v^2 \ln^2(2\sigma)}{2\sigma} + 6 \frac{\Delta^{\frac{1}{2}} vw \ln(2\sigma)}{2\sigma} + 3 \frac{\Delta^{\frac{1}{2}} w^2}{2\sigma} + \\ &\quad \left. + v^3 \ln^3(2\sigma) + 3v^2 w \ln^2(2\sigma) + 2vw^2 \ln(\sigma) + w^3 \right), \end{aligned} \quad (69)$$



with

$$\bar{v} = \frac{1}{2} \sum_{n=0}^{\infty} v_{n+1} \sigma^n, \quad \bar{w} = \frac{1}{2} \sum_{n=0}^{\infty} w_{n+1} \sigma^n. \quad (70)$$

The first term is found to be

$$\frac{\Delta^{\frac{3}{2}}}{(2\sigma)^3} \Big|_R^M = \frac{1}{4} (\nabla_a \nabla_b \nabla_c \Delta^{\frac{1}{2}} - 3 \nabla_a \nabla_b \Delta^{\frac{1}{2}}{}_{;c} + 3 \nabla_a \Delta^{\frac{1}{2}}{}_{;bc} - \Delta^{\frac{1}{2}}{}_{;abc}) \sigma^a \sigma^b \sigma^c \frac{\ln(2M^2 \sigma)}{(2\sigma)^3}. \quad (71)$$

Similarly the second and third terms can be computed:

$$\frac{\Delta v_0 \ln(2\sigma)}{(2\sigma)^2} \Big|_R^M = \frac{1}{4} (\nabla_a v_0 - v_{0;a}) \sigma^a \frac{\Delta \ln(2M^2 \sigma) \ln(\frac{2\sigma}{M^2})}{(2\sigma)^2}, \quad (72)$$

$$\frac{\Delta w_0}{(2\sigma)^2} \Big|_R^M = \frac{1}{2} (\nabla_a w_0 - w_{0;a}) \sigma^a \frac{\Delta \ln(2M^2 \sigma)}{(2\sigma)^2}. \quad (73)$$

Without further knowledge about the Feynman propagator, the β -function can be evaluated to lowest order, leading to the result $\beta_2 = 3$ in this renormalization scheme. The calculation requires the use of the identity $\nabla_a \frac{\Delta \sigma^a}{(2\sigma)^2} = 2\pi^2 \delta(p, p')$.

⁵For the σ calculus see e.g. [Ful89]

CONCLUSIONS

The elegant method of CPT is suited not only for the investigation of renormalizability but also for the performance of perturbative computations. The subtraction procedure on the test functions can be pulled back to the distributions yielding Differential Renormalization. Therefore it is possible to work in the standard integral kernel representation.

As the whole procedure is formulated in configuration space, it can be transferred to distributions on a manifold. This enables one to give compact expressions for the renormalization of quantum fields in curved spacetime, at least in the Euclidean case.

ACKNOWLEDGEMENTS

I am grateful to K. Fredenhagen for many helpful discussions and to Marek Radzikowski and Klaus Bresser for their comments on the manuscript.

REFERENCES

- [BF97] R. Brunetti and K. Fredenhagen, *Interacting quantum fields in curved space: Renormalizability of φ^4* , gr-qc/ **9701048** (1997), To be published in the proceedings of the conference "Operator Algebras and Quantum Field Theory", held at Academia Nazionale dei Lincei, Rome, Italy, July 1996.
- [BFK96] R. Brunetti, K. Fredenhagen, and M. Köhler, *The microlocal spectrum condition and wick polynomials of free fields on curved spacetimes*, Comm. Math. Phys. **180** (1996), 633–652.
- [BS75] Philippe Blanchard and Roland Seneor, *Green's functions in theories with massless particles (in perturbation theory)*, Ann. Inst. H. Poincaré Sect. A **XXIII** (1975), no. 2, 147–209.
- [BS76] N. N. Bogoliubov and D. V. Shirkov, *Introduction to the theory of quantized fields*, 3rd ed., John Wiley and Sons, 1976.
- [Bun81] T. S. Bunch, *BPHZ renormalization of $\lambda\phi^4$ field theory in curved spacetime*, Ann. Physics **131** (1981), 118–148.
- [DB60] Bryce S. DeWitt and Brehme, *Radiation damping in a gravitational field*, Ann. Physics **9** (1960), 220–259.
- [DHKS94a] M. Dütsch, T. Hurth, F. Krahe, and G. Scharf, *Causal construction of Yang-Mills theories.-i*, Nuovo Cimento A **106** (1994), 1029–1041.
- [DHKS94b] M. Dütsch, T. Hurth, F. Krahe, and G. Scharf, *Causal construction of Yang-Mills theories.-ii*, Nuovo Cimento A **107** (1994), 375–406.
- [DHKS95] M. Dütsch, T. Hurth, F. Krahe, and G. Scharf, *Causal construction of Yang-Mills theories.-iii*, Nuovo Cimento A **108** (1995), 679–708.
- [DHS95] M. Dütsch, T. Hurth, and G. Scharf, *Causal construction of Yang-Mills theories.-iv. unitarity*, Nuovo Cimento A **108** (1995), 737–774.
- [DM75] H. G. Dosch and V. F. Müller, *Renormalization of quantum electrodynamics in an arbitrarily strong time independent external field*, Fortschr. Phys. **23** (1975), 661–689.
- [EG73] H. Epstein and V. Glaser, *The role of locality in perturbation theory*, Ann. Inst. H. Poincaré Sect. A **XIX** (1973), no. 3, 211–295.
- [FJL92] Daniel Z. Freedman, Kenneth Johnson, and José I. Latorre, *Differential regularization and renormalization: a new method of calculation in quantum field theory*, Nuclear Phys. B **371** (1992), 352–414.
- [FLS64] Richard P. Feynman, Robert B. Leighton, and Matthew Sands, *Lectures on physics*, Addison-Wesley, 1964.
- [Fre96] K. Fredenhagen, *Renormierungstheorie*, Lecture notes, October 1996.
- [Ful89] S. Fulling, *Aspects of quantum field theory in curved spacetime*, Cambridge University Press, 1989.
- [HL92] Peter E. Haagensen and José I. Latorre, *Differential renormalization of massive quantum field theories*, Phys. Lett. B **283** (1992), 293–297.

- [Hör90] L. Hörmander, *The analysis of linear partial differential operators i*, 2nd ed., Springer, 1990.
- [KW91] B. S. Kay and R. M. Wald, *Theorems on the uniqueness and thermal properties of stationary, nonsingular, quasifree states on spacetimes with a bifurcate killing horizon*, Phys. Rep. **207** (1991), no. 2, 49–136.
- [Lüs82] M. Lüscher, *Dimensional regularisation in the presence of large background fields*, Ann. Physics **142** (1982), 359–392.
- [Sch95] G. Scharf, *Finite quantum electrodynamics*, 2nd ed., Springer, 1995.
- [Ste71] O. Steinmann, *Perturbation expansions in axiomatic field theory*, Lecture Notes in Physics, vol. 11, Springer, 1971.
- [Sto93] R. Stora, *Differential algebras in lagrangean field theory*, ETH Lectures, January-February 1993.
- [SZ93] V. A. Smirnov and O. I. Zav'yalov, *On differential renormalization*, Theoret. and Math. Phys. **96** (1993), 974–981.